

UNIVERSAL BAD NEWS PRINCIPLE AND PRICING OF OPTIONS ON DIVIDEND-PAYING ASSETS

SVETLANA BOYARCHENKO* AND SERGEI LEVENDORSKIĬ†

* Department of Economics, The University of Texas at Austin,
1 University Station C3100, Austin, TX 78712, U.S.A.
e-mail: sboyarch@eco.utexas.edu

† Department of Economics, The University of Texas at Austin,
1 University Station C3100, Austin, TX 78712, U.S.A.
e-mail: leven@eco.utexas.edu

ABSTRACT. We solve the pricing problem for perpetual American puts and calls on dividend-paying assets. The dependence of a dividend process on the underlying stochastic factor is fairly general: any non-decreasing function is admissible. The stochastic factor follows a Lévy process. This specification allows us to consider assets that pay no dividends at all when the level of the underlying factor (say, the assets of the firm) is too low, and assets that pay dividends at a fixed rate when the underlying stochastic process remains in some range. Certain dividend processes exhibiting mean-reverting features can be modelled as appropriate increasing functions of Lévy processes. The pay-offs of both the American put and call options can be represented as the expected present value (EPV) of a certain stream of dividends: $g(X_t) = \delta(X_t) - qK$ and $g(X_t) = qK - \delta(X_t)$, respectively, and we show that the option must be exercised the first time the EPV of the stream $g(\underline{X}_t)$, where $\underline{X}_t = \inf_{0 \leq s \leq t} X_s$ is the infimum process starting from the current level X_0 , becomes positive. Thus, the exercise threshold depends only on the record setting bad news. The results can be applied to the theory of real options as well; as one of possible applications, we consider the problem of incremental capital expansion.

1. INTRODUCTION

The objective of this paper is to provide a general framework for pricing and optimal exercise strategies for American options on dividend-paying assets, for fairly general dependence of dividend rate, $\delta(X_t)$, on the underlying stochastic factor. The stochastic factor, X_t , follows a Lévy process. The standard approach uses the price of an asset as the

primitive; we show that the use of the dividend process as a primitive has certain advantages. In the case of an asset which evolves as a geometric Brownian motion or Lévy process, and pays dividends as a constant proportion of the asset's value, both specifications can easily be transformed one into another. The use of the dividend process as a primitive allows us to consider assets that pay no dividends at all when the level of the underlying factor (say, the value of the firm) is too low, and assets that pay dividends at a fixed rate when the underlying stochastic process remains in some range. Certain dividend processes exhibiting mean-reverting features can be modelled as appropriate increasing functions of Lévy processes. In this paper, we consider perpetual American put and call options; by using the variant of Carr's randomization procedure developed in Levendorskiĭ (2004), which is, essentially, a sequence of embedded perpetual options, it is possible to apply the method of the paper to the case of American options with finite time horizon.

Given a candidate for the optimal exercise threshold, we calculate the option value, and the form of the solution suggests the following description of the optimal exercise strategy. If the payoff stream is a decreasing function of the underlying stochastic factor, then it is optimal to exercise a put-like option the first time the EPV of the stream of payoffs calculated for the supremum process instead of the original stochastic process becomes non-positive. Similarly, if the payoff stream is an increasing function of the underlying stochastic factor, then it is optimal to exercise a call-like option the first time the EPV of the stream of payoffs calculated for the infimum process instead of the original stochastic process becomes non-negative. This allows us to formulate a general optimal exercise rule: it is optimal to exercise the right for (respectively, to give up) the stream of stochastic payoffs, g_t when the EPV of the stream $\underline{g}_t = \inf_{0 \leq s \leq t} g_s$, becomes non-negative (respectively, non-positive). We call the above statement a *universal record setting bad news principle*. This principle naturally generalizes and extends Bernanke's (1983) *bad news principle* and record setting news principles spelled out in Boyarchenko (2004). In the latter paper, the principles were stated and proved for the streams of the form $e^{X_t} - K$ and $K - e^{X_t}$, where X_t is a Lévy process. Here the result is proved for any monotone function, $g_t = g(X_t)$. The method of the paper works for some non-monotone payoff streams as well.

If the underlying process is a diffusion with exponentially distributed jumps, calculation of the optimal exercise price and rational option price reduce to calculation of simple integrals, and solution of one

equation. If, in addition, the dividend process is a piece-wise constant function, or more generally, piece-wise exponential polynomial, then all the integrals can be calculated explicitly, and the optimal exercise price can be found as a unique solution of an algebraic equation with a monotone function.

Now we describe the findings of the paper in more details. Let $\delta(X_t)$ be the dividend process on the asset. The riskless rate $q > 0$ is fixed. Assume that the underlying stochastic factor $\{X_t\}$ is a Lévy process under a risk-neutral measure chosen by the market, denoted $\mathbb{Q}$, and let $(\Omega, \mathcal{F}, \mathbb{Q})$ be the corresponding probability space (for general definitions of the theory of Lévy processes, see, e.g., Bertoin (1996) and Sato (1999)). Then the EPV of the stream $g(X_t)$, at the spot level $X_0 = x$, is given by

$$(1.1) \quad (U_X^q \delta)(x) := E^x \left[\int_0^\infty e^{-qt} \delta(X_t) dt \right].$$

For the perpetual American call on the asset, the payoff function is

$$G(x) = (U_X^q \delta)(x) - K = U_X^q(\delta(\cdot) - qK)(x),$$

where K is the strike price, and for the perpetual American put, the payoff function is

$$G(x) = K - (U_X^q \delta)(x) = U_X^q(qK - \delta(\cdot))(x).$$

(The standard specification is $G(X_t) = e^{X_t} - K$ and $G(X_t) = K - e^{X_t}$, respectively, where X_t is the log-price of the stock.) The rational price of the option with the payoff $G(X_t)$ is given by

$$(1.2) \quad V(x) = \sup E^x[e^{-q\tau} G(X_\tau)],$$

where E^x denotes the expectation under $\mathbb{Q}$, and the supremum is taken over a set $\mathcal{M}$ of all stopping times $\tau = \tau(\omega)$ satisfying $0 \leq \tau(\omega) \leq +\infty$, $\omega \in \Omega$; if $\tau(\omega) = +\infty$, then $G(\tau(\omega)) = 0$ by definition (see, e.g., Shiryaev (1999), XVIII, 2). Notice that we use $G(X_t)$ rather than $\max\{G(X_t), 0\}$, which is admissible because the option is not exercised unless $G(X_t)$ is positive (the equivalence of these two specifications was used in Darling et al. (1972)). In the paper, the optimal stopping time, τ , turns out to be the hitting time of a semi-finite interval of the form $(-\infty, h]$ (put-like options) or $[h, +\infty)$ (call-like options). We denote these hitting times by τ_h^- and τ_h^+ , respectively. The class of the hitting times of semi-finite intervals is denoted by $\mathcal{M}_0$.

Perpetual American options were considered by many authors, both in discrete and continuous time models. Mc Kean (1965) calculated the exercise boundary and price for perpetual call option in the continuous time Gaussian model, Darling et al (1972) solved the corresponding

problem in the discrete time model, for arbitrary random walk, and Merton (1973) solved the problem for the put in the continuous time Gaussian model. Starting from the middle of 1990-th, a series of results for Lévy processes of varying degree of generality were obtained by various authors, using different methods (see the bibliography in Boyarchenko and Levendorskiĭ (2002a, b) and Mordecki (2002), and a more detailed discussion in Section 5.

In the current paper, we present the solution to the optimal stopping problem for wide class of Lévy processes satisfying the (ACP)–property (absolute continuity of potential kernels: see, e.g., Sato (1999)), and fairly general payoff functions. We formulate our results in terms of a stream of payoffs, g (dividends), whose expected present value (EPV) is equal to the given payoff, G (spot-price of the stock), and *expected present value operators*, U_X^q and $U_{\underline{X}}^q$, of the supremum and infimum processes (these operators are defined by formulas similar to (1.1); for details, see Section 3). In the case of the put on a stock which pays no dividends and similar put-like options, we formulate the results separately, in terms of the payoff function itself. The reason is that the price of a stock which pays no dividends cannot be determined as the EPV of any stream. For the case of call-like options (the case of an increasing G), we prove the optimality in the class $\mathcal{M}$, under the weak conditions that the EPV of the stream g under the infimum process, $U_{\underline{X}}^q g$, changes sign from “-” to “+” only once, and the stream g is a non-decreasing function. The last condition is not necessary; in fact, the proof in the paper works in some situations when the stream is not monotone. Similar results are proved for put-like options with decreasing payoff functions G . This time, the optimal exercise price is the zero of the EPV of the stream g under the supremum process, $U_X^q g$, which is assumed to change sign from “+” to “-” only once.

The rest of the paper is organized as follows. In Section 2, we recall the basic definitions of the theory of Lévy processes, introduce the EPV operators of the supremum and infimum processes, and calculate their action for the case of diffusions with exponentially distributed jumps. In Section 3 formulate the main results, and explicitly calculate the optimal exercise thresholds and option prices for diffusions with exponentially distributed jumps. The proofs of the main results are in Section 4–Section 6. In Section 7, we apply the method of the paper to the problem of capital expansion.

2. LÉVY PROCESSES

2.1. Recall that a Lévy process is a process with stationary independent increments (for general definitions, see e.g. Sato (1999)). A Lévy process may have a Gaussian component and/or pure jump component. The latter is characterized by the density of jumps, which is called the Lévy density. We denote it by $F(dx)$. Also, a Lévy process can be completely specified by its Lévy exponent, Ψ , definable from the moment-generating function $E[e^{zX_t}] = e^{t\Psi(z)}$ (we confine ourselves to the one-dimensional case). If X_t is a Lévy process with finite variation jump component, then the Lévy exponent is given by

$$(2.1) \quad \Psi(z) = bz + \frac{\sigma^2}{2}z^2 + \int_{-\infty}^{+\infty} (e^{zy} - 1)F(dy),$$

where σ^2 and b are the variance and drift coefficient of the Gaussian component, and $F(dy)$ satisfies

$$\int_{\mathbb{R} \setminus \{0\}} \min\{1, |y|\} F(dy) < +\infty.$$

Equation (2.1) is a special case of the Lévy-Khintchine formula; for the general case, see e.g. Sato (1999). In this paper, we will illustrate our general results for the case of the Lévy density

$$(2.2) \quad F(dx) = c^+ \lambda^+ e^{-\lambda^+ x} \mathbf{1}_{(0, +\infty)}(x) dx + c^- (-\lambda^-) e^{-\lambda^- x} \mathbf{1}_{(-\infty, 0)}(x) dx,$$

where $\lambda^+ > 0 > \lambda^-$, and $c_{\pm} > 0$. Then

$$(2.3) \quad \Psi(z) = \frac{\sigma^2}{2}z^2 + bz + \frac{c^+ z}{\lambda^+ - z} + \frac{c^- z}{\lambda^- - z},$$

where $\sigma^2 > 0$ and $b \in \mathbb{R}$ are the variance and drift of the Gaussian component. The $\Psi(z)$ is analytic in the strip $\Re z \in (\lambda^-, \lambda^+)$.

2.2. The Lévy exponent appears when we calculate the action of the infinitesimal generator of X_t , denoted L , on exponential functions: $Le^{zx} = \Psi(z)e^{zx}$. The EPV-operator (1.1) calculates the expected present value of the dividend stream δ . From the fundamental relation between the infinitesimal generator and the resolvent,

$$(2.4) \quad (q - L)U_X^q = U_X^q(q - L) = I,$$

one concludes that U_X^q acts on exponential functions as the multiplication operator by the number $(q - \Psi(z))^{-1}$:

$$(2.5) \quad U_X^q e^{zx} = (q - \Psi(z))^{-1} e^{zx}.$$

Let $\bar{X}_t = \sup_{0 \leq s \leq t} X_s$ and $\underline{X}_t = \inf_{0 \leq s \leq t} X_s$ be the supremum and infimum processes of X_t . Introduce EPV-operators $U_{\bar{X}}^q$ and $U_{\underline{X}}^q$ of the supremum and infimum processes by

$$U_{\bar{X}}^q g(x) := E^x \left[\int_0^\infty e^{-qt} g(\bar{X}_t) dt \right] := E \left[\int_0^\infty e^{-qt} g(\bar{X}_t) dt \mid X_0 = x \right]$$

and

$$U_{\underline{X}}^q g(x) := E^x \left[\int_0^\infty e^{-qt} g(\underline{X}_t) dt \right] := E \left[\int_0^\infty e^{-qt} g(\underline{X}_t) dt \mid X_0 = x \right],$$

respectively. It is straightforward to check that $qU_{\bar{X}}^q$ and $qU_{\underline{X}}^q$ also act on an exponential function e^{zx} as multiplication operators by certain numbers, which we denote $\kappa_q^+(z)$ and $\kappa_q^-(z)$, respectively:

$$(2.6) \quad qU_{\bar{X}}^q e^{zx} = \kappa_q^+(z) e^{zx}, \quad qU_{\underline{X}}^q e^{zx} = \kappa_q^-(z) e^{zx}.$$

These numbers are

$$(2.7) \quad \kappa_q^+(z) = qE \left[\int_0^\infty e^{-qt} e^{z\bar{X}_t} dt \right],$$

$$(2.8) \quad \kappa_q^-(z) = qE \left[\int_0^\infty e^{-qt} e^{z\underline{X}_t} dt \right].$$

Notice that $\kappa_q^+(z)$ (resp., $\kappa_q^-(z)$) is analytic on the half-plane $\Re z < 0$ (resp., $\Re z > 0$), and continuous up to the boundary. The Wiener-Hopf factorization formula reads (see, e.g., Sato (1999), Section 45)

$$(2.9) \quad \frac{q}{q - \Psi(z)} = \kappa_q^+(z) \kappa_q^-(z).$$

By applying $U_{\bar{X}}^q$, $U_{\underline{X}}^q$ and $U_{\underline{X}}^q$ to an exponential function $g(x) = e^{zx}$ and using (2.5) and (2.6)–(2.9), we obtain

$$(2.10) \quad U_{\bar{X}}^q g(x) = qU_{\bar{X}}^q U_{\underline{X}}^q g(x) = qU_{\underline{X}}^q U_{\bar{X}}^q g(x).$$

By linearity, (2.10) holds for linear combinations of exponents, and integrals of exponents, hence, for wide classes of functions.

Equation (2.10) means that the EPV-operator of a Lévy process admits a factorization into a product of the EPV-operator of the supremum process and the one of the infimum process.

2.3. For diffusions with exponentially distributed jumps, $q - \Psi(z)$ is the rational function, which has 4 real zeroes; two of them are positive, and two negative. We will call them β_j^- and β_j^+ , $j = 1, 2$, respectively. It is easy to show that λ^- separates the negative roots, and λ^+ – the positive ones. We have $\beta_2^- < \lambda^- < \beta_1^+ < 0 < \beta_1^- < \lambda^+ < \beta_2^+$. Since $q - \Psi(z)$ is rational, the factors $\kappa^\pm(z)$ can easily be obtained by

representing the LHS in (2.9) as the fraction of two polynomials, factorizing these polynomials out, and collecting the factors with positive (respectively, negative) zeroes. For details of these calculations and calculations below, see Levendorskiĭ (2004). We obtain

$$(2.11) \quad \kappa_q^+(z) = \frac{\beta_1^+ \beta_2^+ (\lambda^+ - z)}{(\beta_1^+ - z)(\beta_2^+ - z)\lambda^+}$$

$$(2.12) \quad = \sum_{j=1,2} \frac{a_j^+}{\beta_j^+ - z},$$

where

$$(2.13) \quad a_1^+ = \frac{\beta_1^+ \beta_2^+ (\lambda^+ - \beta_1^+)}{(\beta_2^+ - \beta_1^+)\lambda^+}, \quad a_2^+ = \frac{\beta_1^+ \beta_2^+ (\lambda^+ - \beta_2^+)}{(\beta_1^+ - \beta_2^+)\lambda^+},$$

are positive, and

$$(2.14) \quad \kappa_q^-(z) = \frac{\beta_1^- \beta_2^- (\lambda^- - z)}{(\beta_1^- - z)(\beta_2^- - z)\lambda^-}$$

$$(2.15) \quad = \sum_{j=1,2} \frac{a_j^-}{\beta_j^- - z},$$

where

$$(2.16) \quad a_1^- = \frac{\beta_1^- \beta_2^- (\lambda^- - \beta_1^-)}{(\beta_2^- - \beta_1^-)\lambda^-}, \quad a_2^- = \frac{\beta_1^- \beta_2^- (\lambda^- - \beta_2^-)}{(\beta_1^- - \beta_2^-)\lambda^-}$$

are negative. The operators $qU_{\bar{X}}^q$ and $qU_{\underline{X}}^q$ act as follows:

$$(2.17) \quad qU_{\bar{X}}^q u(x) = \sum_{j=1,2} a_j^+ \int_0^\infty e^{-\beta_j^+ y} u(x+y) dy,$$

$$(2.18) \quad qU_{\underline{X}}^q u(x) = \sum_{j=1,2} (-a_j^-) \int_{-\infty}^0 e^{-\beta_j^+ y} u(x+y) dy.$$

To see this, it suffices to insert $u(x) = e^{zx}$, and use the definition of the numbers $\kappa_q^+(z), \kappa_q^-(z)$.

2.4. In applications, one needs to calculate the EPV's of exponentially growing payoffs. To ensure that such EPV's are finite, Ψ must be defined not only on the imaginary axis $i\mathbb{R}$ but on a strip of the form $\lambda^- \leq \Re z \leq \lambda^+$, where $\lambda^- \leq 0 < \lambda^+$. An equivalent condition is $\Psi(z) < +\infty$, for all $z \in [\lambda^-, \lambda^+]$. Since $\Psi(0) = 0$ and $q > 0$, there exist $\lambda^- \leq \sigma_- \leq 0 < \sigma_+ \leq \lambda^+$ such that

$$(2.19) \quad q - \Psi(z) > 0, \quad \sigma_- \leq z \leq \sigma_+.$$

In this case, $q - \Psi(z)$ does not vanish in the strip $\Re z \in [\sigma_-, \sigma_+]$, the domains of the definition of κ_q^+ and κ_q^- contain this strip, and the equality (2.9) holds for z in the strip $\Re z \in [\sigma_-, \sigma_+]$.

3. MAIN RESULTS

3.1. Assume that the dividend stream is a non-negative non-decreasing piece-wise continuous function of the stochastic factor X_t , which admits a bound

$$(3.1) \quad \delta(x) \leq C(1 + e^{\sigma_+ x}).$$

Example 3.1. The asset pays no dividends when the underlying stochastic factor (say, value of the firm) is too low: $\delta(x) = 0$, $x \leq x_0$, and the dividends increase exponentially after the factor crosses a certain level:

$$(3.2) \quad \delta(x) = (e^x - e^{x_0})^+ := \max\{0, e^x - e^{x_0}\}.$$

Example 3.2. The asset pays no dividends when the underlying stochastic factor (say, value of the firm) is too low: $\delta(x) = 0$, $x \leq x_0$, and the dividends increase not so fast after the factor crosses a certain level:

$$(3.3) \quad \delta(x) = (x - x_0)^+.$$

Example 3.3. The asset pays no dividends when the underlying stochastic factor (say, value of the firm) is too low: $\delta(x) = 0$, $x \leq x_0$. When the critical level is crossed, the dividends increase but eventually the growth slows and essentially stops:

$$(3.4) \quad \delta(x) = (1 - e^{-x})^+.$$

Example 3.4. The asset pays dividends at a fixed rate when the underlying process is within a certain range; when the process arrives in the next range, the dividend rate changes by a jump:

$$(3.5) \quad \delta(x) = \sum_j \delta_j \mathbf{1}_{[d_j, +\infty)}(x).$$

The sum can be finite (in this case, the dividends are capped, as in Example 3.3) or infinite, which allows for unbounded growth of dividends.

Example 3.5. The dividends are paid in the constant proportion to the firm's value: $d(X_t) = dX_t$, but the value itself is an increasing function of a Lévy process, Y_t : $X_t = f(Y_t)$. If $f(y)$ is convex for $y < y_0$, and concave for $y > y_0$, the process X_t may exhibit a mean-reverting feature.

Example 3.6. We can easily generalize Examples 3.1–3.4 by using $\delta(f(Y_t))$, where f is an increasing function, and Y_t is a Lévy process.

In the following sections, we prove the following results for wide classes of Lévy processes, including diffusions with exponentially distributed jumps.

3.2. First, we consider the perpetual American call. Let h^* be the solution to the equation

$$(3.6) \quad w(x) := (U_{\underline{X}}^q \delta)(x) - K = 0.$$

(The solution exists if $\delta(x)$ is sufficiently large for large x , and it is unique since δ is monotone). Then h^* is the optimal exercise level for the perpetual American call on the asset with the dividend stream δ . After h^* is found, we calculate the rational call price, for $x \leq h_*$:

$$(3.7) \quad V^*(x) = (qU_{\bar{X}}^q \mathbf{1}_{[h^*, +\infty)} w)(x).$$

For exponential jump-diffusions, we use (2.18) and rewrite (3.6) as

$$(3.8) \quad q^{-1} \sum_{j=1,2} (-a_j^-) \int_{-\infty}^0 e^{-\beta_j^- y} \delta(x+y) dy - K = 0.$$

The solution to (3.8) can easily be found by standard numerical methods. In particular, in Examples 3.1–3.4 (and in many others), the integral in (3.12) can be calculated explicitly, and we have to solve an algebraic equation. Consider, for instance, Example 3.1. Without loss of generality, set $x_0 = 0$. Then for $x \leq 0$, the LHS is $-K$, therefore the root is on the positive half-axis. For $x > 0$, we change the variables

$$(3.9) \quad q^{-1} \sum_{j=1,2} (-a_j^-) e^{\beta_j^- x} \int_{-\infty}^x e^{-\beta_j^- y} \delta(y) dy - K = 0,$$

and calculate

$$\int_{-\infty}^x e^{-\beta_j^- y} \delta(y) dy = \int_0^x e^{-\beta_j^- y} (e^y - 1) dy = \frac{e^{(1-\beta_j^-)x} - 1}{1 - \beta_j^-} - \frac{e^{-\beta_j^- x} - 1}{-\beta_j^-}.$$

We see that the first term on the RHS of (3.9) is

$$\begin{aligned}
& q^{-1} \sum_{j=1,2} (-a_j^-) \left\{ \frac{e^x - e^{-\beta_j^- x}}{1 - \beta_j^-} - \frac{1 - e^{-\beta_j^- x}}{-\beta_j^-} \right\} \\
&= q^{-1} \sum_{j=1,2} \left\{ \frac{-a_j^-}{1 - \beta_j^-} e^x - \frac{a_j^-}{\beta_j^-} + \frac{a_j^- e^{-\beta_j^- x}}{(1 - \beta_j^-) \beta_j^-} \right\} \\
&= q^{-1} \left\{ \kappa_q^-(1) e^x - 1 + \sum_{j=1,2} \frac{a_j^-}{(1 - \beta_j^-) \beta_j^-} e^{-\beta_j^- x} \right\}.
\end{aligned}$$

Thus, the equation for h^* is

$$\kappa_q^-(1) e^x - 1 + \sum_{j=1,2} \frac{a_j^-}{(1 - \beta_j^-) \beta_j^-} e^{-\beta_j^- x} = qK + 1, \quad x > 0,$$

and it can be easily solved. The equation being solved, we represent w in (3.6) in the form

$$w(x) = q^{-1} \sum_{j=1,2} (-a_j^-) \int_{-\infty}^0 e^{-\beta_j^- y} (\delta(x+y) - \delta(h_* + y)) dy,$$

and calculate the rational call price using (2.17):

$$\begin{aligned}
V^*(x) &= \sum_{k=1,2} a_k^+ \int_0^{+\infty} e^{-\beta_k^+ y} \mathbf{1}_{[h^*, +\infty)}(x+y) w(x+y) dy \\
&= \sum_{k=1,2} a_k^+ \int_{h^*-x}^{+\infty} e^{-\beta_k^+ y} w(x+y) dy \\
&= q^{-1} \sum_{j,k=1,2} a_k^+ (-a_j^-) \\
&\quad \cdot \int_{h^*-x}^{+\infty} \int_{-\infty}^0 e^{-\beta_k^+ y - \beta_j^- z} [\delta(x+y+z) - \delta(h^* + z)] dz dy \\
&= q^{-1} \sum_{j,k=1,2} a_k^+ (-a_j^-) e^{\beta_k^+(x-h^*)} \\
&\quad \cdot \int_0^{+\infty} \int_{-\infty}^0 e^{-\beta_j^- z - \beta_k^+ y} [\delta(h^* + y+z) - \delta(h^* + z)] dz dy \\
&= q^{-1} \sum_{k=1,2} a_k^+ \mathcal{D}_k^+ e^{\beta_k^+(x-h^*)},
\end{aligned}$$

where the constants

$$\mathcal{D}_k^+ = \sum_{j=1,2} (-a_j^-) \int_0^{+\infty} \int_0^{+\infty} e^{\beta_j^- z - \beta_k^+ y} [\delta(h^* - z) - \delta(h^* + y - z)] dz dy$$

can be calculated quite easily in all Examples 3.1-3.4 (and in many other examples).

3.3. Now we consider the put. Let h_* be the solution to the equation

$$(3.10) \quad w(x) := K - (U_X^q \delta)(x) = 0.$$

(The solution exists, if $\delta(x) = 0$ or sufficiently small for x in a neighborhood of $-\infty$, and it is unique, since δ is non-decreasing). Then h_* is the optimal exercise level for the perpetual American put on the asset with the dividend stream δ . After h_* is found, we calculate the rational put price, for $x \geq h_*$:

$$(3.11) \quad V_*(x) = (qU_X^q \mathbf{1}_{(-\infty, h_*]} w)(x).$$

For exponential jump-diffusions, we use (2.17) and rewrite (3.10) as

$$(3.12) \quad K - q^{-1} \sum_{k=1,2} a_k^+ \int_0^{+\infty} e^{-\beta_k^+ y} \delta(x+y) dy = 0.$$

The solution to (3.12) can easily be found by standard numerical methods. In particular, in Examples 3.1-3.4 (and in many others), the integral in (3.12) can be calculated explicitly, and we have to solve an algebraic equation. The equation being solved, we represent w in (3.10) in the form

$$w(x) = q^{-1} \sum_{k=1,2} a_k^+ \int_0^{+\infty} e^{-\beta_k^+ y} (\delta(h_* + y) - \delta(x+y)) dy,$$

and calculate the rational put price using (2.18):

$$\begin{aligned}
V_*(x) &= \sum_{j=1,2} (-a_j^-) \int_{-\infty}^0 e^{-\beta_j^- y} \mathbf{1}_{(-\infty, h_*]}(x+y) w(x+y) dy \\
&= \sum_{j=1,2} (-a_j^-) \int_{-\infty}^{h_*-x} e^{-\beta_j^- y} w(x+y) dy \\
&= q^{-1} \sum_{j,k=1,2} a_k^+ (-a_j^-) \\
&\quad \cdot \int_{-\infty}^{h_*-x} \int_0^{+\infty} e^{-\beta_j^- y - \beta_k^+ z} [\delta(h_* + z) - \delta(x+y+z)] dz dy \\
&= q^{-1} \sum_{j,k=1,2} a_k^+ (-a_j^-) e^{\beta_j^- (x-h_*)} \\
&\quad \cdot \int_{-\infty}^0 \int_0^{+\infty} e^{-\beta_j^- y - \beta_k^+ z} [\delta(h_* + z) - \delta(h_* + y + z)] dz dy \\
&= q^{-1} \sum_{j=1,2} (-a_j^-) \mathcal{D}_j^- e^{\beta_j^- (x-h_*)},
\end{aligned}$$

where the constants

$$\mathcal{D}_j^- = \sum_{k=1,2} a_k^+ \int_0^{+\infty} \int_0^{+\infty} e^{\beta_j^- y - \beta_k^+ z} [\delta(h_* + z) - \delta(h_* - y + z)] dz dy$$

can be calculated quite easily in Examples 3.1-3.4 (and in many other examples).

4. THE WIENER-HOPF METHOD AND SOME APPLICATIONS

4.1. From now on, our standing assumption is that the Lévy process X satisfies the (ACP)-property (for the definition, see, e.g., Bertoin (1996) and Sato (1999)). Fix $h \in \mathbb{R}$, and set $\tau = \tau_h^- = \inf\{t \mid X_t < h\}$. For z in the upper right-plane $\Re z \geq 0$, consider functions

$$f(z; x) = f(q, h, z; x) = E^x[\exp(zX_\tau - q\tau)],$$

and

$$f^1(z; x) = f^1(q, h, z; x) = \kappa_q^-(z)^{-1} q U_{\underline{X}}^q u(z; x),$$

where $u(z; x) = u(h, z; x) = \mathbf{1}_{(-\infty, h]}(x) e^{zx}$.

Lemma 4.1. *For $x > h$, $f(z; x) = f^1(z; x)$.*

Proof. For a fixed x , both functions are analytic in the half-plane $\Re z > 0$, and continuous up to the boundary. Hence, it suffices to prove the

equality for $z \in [0, \epsilon]$, where ϵ is some positive number. If $z = 0$, then $\kappa_q^-(z)^{-1} = 1$, and the equality

$$E^x[\exp(zX_\tau - q\tau)] = qE^x \left[\int_0^{+\infty} e^{-qt} \mathbf{1}_{(-\infty, h]}(\underline{X}_t) dt \right]$$

holds. Thus, the lemma is proved in the case $z = 0$. The proof for small positive z consists of the following steps: $f(z; \cdot)$ is RCLL on $(h; +\infty)$ (right continuous with left limits); $f^1(z; \cdot)$ is RCLL on $(h; +\infty)$; the Laplace transforms of these two functions are equal.

Function $f(0; \cdot)$ is q -excessive (Proposition 41.5 (ii) and (viii) in Sato (1999)). Since X satisfies the (ACP)-property, a q -excessive function is lower semi-continuous (Theorem 41.5 (4) in Sato (1999)), but $f(0; \cdot)$ is evidently non-increasing; hence, $f(0; \cdot)$ and $f^1(0; \cdot)$ are RCLL on $(h, +\infty)$. Consider sufficiently small $z > 0$ so that $q - \Psi(z) > 0$. Introduce $\Psi_z(w) := \Psi(w + z) - \Psi(z)$. This is the Lévy exponent of X_t under the Esscher transform of the measure $\mathbb{Q}$; denote the Esscher transform by $\mathbb{Q}_z$. Let $E^{\mathbb{Q}}$ and $E^{\mathbb{Q}_z}$ be the expectation operators under $\mathbb{Q}$ and $\mathbb{Q}_z$, respectively. We have

$$f(z; x) := E^{\mathbb{Q}; x}[\exp(-q\tau_h^- + zX_{\tau_h^-})] = e^{zx} E^{\mathbb{Q}_z; 0}[\exp(-(q - \Psi(z))\tau_h^-)].$$

Since X satisfies the (ACP)-property under $\mathbb{Q}$, it satisfies the (ACP) property under $\mathbb{Q}_z$. Hence, the last factor on the RHS is RCLL on $(h, +\infty)$, and $f(z; \cdot)$, its product with a continuous function, is RCLL on $(h, +\infty)$ as well.

To prove that $f^1(z; \cdot)$ is RCLL on $(h, +\infty)$, we change the variables $x \mapsto x + h$, so that h becomes 0, and represent $f^1(z; \cdot)$ in the form

$$(4.1) \quad f^1(z; x) = \kappa_q^-(z)^{-1} \left(qU_{\underline{X}}^q \mathbf{1}_{(-\infty, 0]}(x) + qU_{\underline{X}}^q \mathbf{1}_{(-\infty, 0]}(e^{zx} - 1)(x) \right),$$

The function $\mathbf{1}_{(-\infty, 0]}(x)(e^{zx} - 1)$ is continuous, therefore the second term in the brackets on the RHS in (4.1) is continuous. The first term in the brackets equals $f(0; x)$, hence it is RCLL on $(h, +\infty)$. We conclude that the sum is RCLL on $(h, +\infty)$.

Now we consider the Laplace transforms. The fluctuation identity (3.13) in Hilberink–Rogers (2002) states that for any $\mu > z$,

$$\int_0^\infty e^{-\mu x} f(z; x) dx = \frac{1}{\mu - z} \left[1 - \frac{\kappa_q^-(\mu)}{\kappa_q^-(z)} \right],$$

and therefore we need to prove the same equality for f^1 . We have

$$f^1(z; x) = \kappa_q^-(z)^{-1} qU_{\underline{X}}^q (1 - \mathbf{1}_{(0, +\infty)}(x)) e^{zx} = e^{zx} - \kappa_q^-(z)^{-1} qU_{\underline{X}}^q v(z; x),$$

where $v(z; x) = \mathbf{1}_{(0, +\infty)}(x)e^{zx}$. For w in the upper half-plane $\Re w > \Re z$, the Laplace transform

$$\hat{v}(z; w) = \int_0^{+\infty} e^{-wx} v(z; x) dx = \int_0^{+\infty} e^{-x(w-z)} dx = (w - z)^{-1}$$

is well-defined; therefore, using the inverse Laplace transform, we obtain, for any $\sigma > \Re z$,

$$f^1(z; x) = e^{zx} - (2\pi i)^{-1} \int_{-i\infty+\sigma}^{+i\infty+\sigma} e^{wx} \kappa_q^-(z)^{-1} \kappa_q^-(w) (w - z)^{-1} dw.$$

We take $\mu > z$ (and $\sigma \in (z, \mu)$), and calculate the Laplace transform

$$\int_0^{+\infty} e^{-\mu x} f^1(z; x) dx = \frac{1}{\mu - z} - \frac{\kappa_q^-(\mu)}{\kappa_q^-(z)(\mu - z)} = \frac{1}{\mu - z} \left[1 - \frac{\kappa_q^-(\mu)}{\kappa_q^-(z)} \right].$$

Thus, the Laplace transforms of $f(z; \cdot)$ and $f^1(z; \cdot)$ are equal, which completes the proof. $\square$

Below, we consider piece-wise continuous streams g ; this condition can be relaxed.

Theorem 4.2. *Assume that the infimum process is non-trivial, and g is non-negative. Then*

$$(4.2) \quad E^x \left[e^{-q\tau_h^-} U_X^q g(X_{\tau_h^-}) \right] = q U_X^q \mathbf{1}_{(-\infty, h]} U_X^q g(x), \quad \forall x > h.$$

Proof. Consider first $g(x) = e^{zx}$, where $z \in i\mathbb{R}$. Using (2.5) and (2.6), we can rewrite (4.2) as

$$(q - \Psi(z))^{-1} f(z; x) = q^{-1} \kappa_q^+(z) f^1(z; x),$$

where f and f^1 are the functions in Lemma 4.1. This equality holds on the strength of (2.10) and Lemma 4.1, hence (4.2) is proved for oscillating exponents.

Next, consider $g \in C_0^\infty(\mathbb{R})$. We represent g as the Fourier integral, use Lemma 4.1 under the integral sign, and obtain (4.2). Finally, a general piece-wise continuous g can be approximated by a sequence $\{g_n\} \in C_0^\infty(\mathbb{R})$, which converges to g point-wise, from below. For each n , (4.2) with g_n instead of g holds. By the Monotone Convergence Theorem, the LHS and RHS with g_n instead of g have point-wise limits, which are the LHS and RHS with g . $\square$

4.2. Fix $h \in \mathbb{R}$, and set $\tau = \tau_h^+ = \inf\{t \mid X_t > h\}$. If we change the direction of the real line, a neighborhood of $-\infty$ becomes a neighborhood of $+\infty$, the supremum process becomes the infimum process, and vice versa. Hence, by symmetry, we obtain

Theorem 4.3. *Assume that the supremum process is non-trivial, and g is non-negative. Then*

$$(4.3) \quad E^x \left[e^{-q\tau_h^+} U_X^q g(X_{\tau_h^+}) \right] = q U_X^q \mathbf{1}_{[h, +\infty)} U_X^q g(x), \quad \forall x < h.$$

4.3. Let (2.19) hold. If g is piece-wise continuous and satisfies an estimate

$$(4.4) \quad |g(x)| \leq C e^{\sigma_- x}, \quad x < h,$$

then the LHS in (4.2) is finite, and (4.2) holds. Similarly, if g is piece-wise continuous and

$$(4.5) \quad |g(x)| \leq C e^{\sigma_+ x}, \quad x > h,$$

then the LHS in (4.3) is finite, and (4.3) holds.

5. OPTIMAL EXERCISE BOUNDARY AND RATIONAL PRICE OF A PERPETUAL PUT-LIKE OPTION

5.1. We use the following general lemma (Lemma 5.1 in Boyarchenko and Levendorskiĭ (2002b) and Lemma 7.1 in Boyarchenko and Levendorskiĭ (2002b)). We formulate it for a special case of a process on the line, and an optimal stopping region of the form $(-\infty, h]$.

Lemma 5.1. *Let h_* and a function V_* satisfy the following conditions:*

$$(5.1) \quad (q - L)V_*(x) = 0, \quad x > h_*;$$

$$(5.2) \quad V_*(x) = G(x), \quad x \leq h_*;$$

$$(5.3) \quad V_*(x) \geq \max\{G(x), 0\}, \quad x \in \mathbb{R};$$

$$(5.4) \quad (q - L)V_*(x) \geq 0, \quad x < h_*;$$

$$(5.5) \quad W_* := (q - L)V_* \text{ is universally measurable;}$$

$$(5.6) \quad U_X^q W_* = V_*.$$

Then the pair $(\tau_{h_*}^-, V_*)$ is the solution to the optimal stopping problem (1.2) in the class $\mathcal{M}$.

5.2. The optimal stopping problem for a put-like option is trivial if the infimum process is trivial. Hence, we presume in this section that the infimum process is non-trivial.

Fix arbitrary h . If X satisfies the (ACP)-property, and G is continuous and does not grow too fast at infinity, so that $V_0^-(h; x) = E^x \left[e^{-q\tau_h^-} G(X_{\tau_h^-}) \right]$ is finite (a sufficient condition is (4.4)), then $V_0^-(h; x)$ satisfies (5.1) with h instead of h_* (see Theorem 2.12 and Remark 2.1 in Boyarchenko and Levendorskiĭ (2002b) or Theorem 2.1 in Boyarchenko and Levendorskiĭ (2002c)), and clearly, $V_0^-(h; x)$ satisfies (5.2) for $x < h$. By Theorem 4.2, for $x > h$, $V_0^-(h; x) = V^-(h; x) := qU_X^q \mathbf{1}_{(-\infty, h]} U_X^q g(x)$. To prove the equality $V_0^-(h; x) = V^-(h; x)$ for $x < h$, we use (2.10) to represent $V^-(h; x)$ in the form

$$\begin{aligned} V^-(h; x) &= qU_X^q U_X^q g(x) - q(U_X^q \mathbf{1}_{(h, +\infty)} w)(x) \\ &= U_X^q g(x) - v(h; x) \\ (5.7) \quad &= G(x) - v(h; x), \end{aligned}$$

where $v(h; x) = q(U_X^q \mathbf{1}_{(h, +\infty)} w)(x)$. For $x \leq h$,

$$v(h; x) = qE \left[\int_0^{+\infty} e^{-qt} (\mathbf{1}_{(h, +\infty)} w)(\underline{X}_t) \mid X_0 = x \right] = 0,$$

therefore $V^-(h; x) = G(x)$, $x \leq h$. But for $x < h$, $E^x \left[e^{-q\tau_h^-} U_X^q g(X_{\tau_h^-}) \right] = G(x)$ as well.

It remains to find h_* such that $V^-(h; x)$ is continuous, and satisfies (5.3)–(5.6). Assume that g is piece-wise continuous, and satisfies (4.4). X satisfies the (ACP)-property, hence $G = U_X^q g$ is continuous, and G satisfies (4.4), since g does. Introduce $w = U_X^q g$, and assume that w is a continuous function that satisfies

$$(5.8) \quad w \text{ changes sign from “+” to “−”, and only once.}$$

If g is continuous, w is continuous as well, and a sufficient condition for (5.8) is that g is decreasing. This condition makes a perfect economic sense for a perpetual put-like option: the stream of payoffs increases when the stochastic factor decreases.

Denote by h_* the solution of the equation

$$(5.9) \quad w(x) = U_X^q g(x) = 0,$$

and set $V_*(\cdot) = V^-(h_*; \cdot)$.

Theorem 5.2. *Let (5.8) hold, and g be non-increasing. Then $(\tau_{h_*}^-, V_*)$ is the optimal solution to the stopping problem, in the class $\mathcal{M}$.*

Proof. Due to the choice of h_* , $\mathbf{1}_{(-\infty, h_*]} w$ is continuous and non-negative. Therefore, $V^-(h^*; \cdot)$ is continuous, and

$$\begin{aligned} V^-(h_*; x) &= qU_{\underline{X}}^q \mathbf{1}_{(-\infty, h_*]} w(x) \\ (5.10) \quad &= qE \left[\int_0^{+\infty} e^{-qt} (\mathbf{1}_{(-\infty, h_*]} w)(x + \underline{X}_t) dt \right] \geq 0, \quad \forall x. \end{aligned}$$

Further, consider (5.7) with $h = h_*$, the solution to (5.9). Due to (5.8), $\mathbf{1}_{(h_*, +\infty)} w$ is non-positive, hence $v(h_*; \cdot)$ in (5.7) is non-positive, and we conclude from (5.7) that $V^-(h_*; x) \geq G(x)$ for all x . Thus, V_* satisfies (5.3). Since (5.1) holds, we need to check (5.4) on $(-\infty, h_*)$. Below, we will show that

$$(5.11) \quad W_* = (q - L)V_* \quad \text{is non-increasing on } (-\infty, h_*),$$

and

$$(5.12) \quad W_*(h_* - 0) \geq 0.$$

Conditions (5.4) and (5.5) follow immediately from (5.11) and (5.12), and it remains to check (5.6). Since $\mathbf{1}_{(-\infty, h_*]} U_{\tilde{X}}^q g$ is continuous, $V_* = V^-(h_*; \cdot)$ given by the LHS in (4.2) is continuous, and since W_* is universally measurable and X satisfies the (ACP)-property, $U_X^q W_*$ is continuous. Therefore it suffices to prove (5.6) in the sense of generalized functions:

$$\int_{-\infty}^{+\infty} U_X^q W_*(x) u(x) dx = \int_{-\infty}^{+\infty} V_*(x) u(x) dx, \quad \forall u \in C_0^\infty(\mathbb{R}).$$

By the standard duality argument,

$$\int_{-\infty}^{+\infty} U_X^q (q - L)V_*(x) u(x) dx = \int_{-\infty}^{+\infty} V_*(x) (q - \tilde{L})U_{\tilde{X}}^q u(x) dx,$$

where $\tilde{X}$ is the dual process and $\tilde{L}$ its generator. Since $u \in C_0^\infty(\mathbb{R})$, $(q - \tilde{L})U_{\tilde{X}}^q u = u$, and the proof of (5.6) is finished.

It remains to prove (5.11) and (5.12). Represent W_* in the form

$$W_* = (q - L)G - (q - L)w^+,$$

where $w^+ = qU_{\underline{X}}^q \mathbf{1}_{(h_*, +\infty)} w$. Since $\text{supp } w^+ \subset [h_*, +\infty)$, and the Gaussian part of the infinitesimal generator is a local (differential) operator,

we have for $x < h_*$

$$\begin{aligned} W_*(x) &= (q - L)G(x) \\ &\quad + \int_{-\infty}^{+\infty} (w^+(x + y) - w^+(x) - \mathbf{1}_{[-1,1]}(y)(w^+)'(x)) F(dy) \\ &= g(x) + \int_{h_* - x}^{+\infty} w(x + y)F(dy), \end{aligned}$$

where $F(dy)$ is the Lévy density. Since $(q - L)G = g$ is non-increasing, $w = U_{\bar{X}}^q g$ is non-increasing as well, hence both terms on the RHS are non-increasing, and (5.11) is proved.

Finally, assume that (5.12) fails. On the strength of (5.11), W_* must be negative on some interval (h, h_*) , where $h < h_*$. By applying $U_{\bar{X}}^q$ to $W_* = (q - L)V_*$ and using (4.2) and (2.10), we obtain

$$U_{\bar{X}}^q W_* = U_{\bar{X}}^q (q - L)qU_{\bar{X}}^q \mathbf{1}_{(-\infty, h_*]} w = \mathbf{1}_{(-\infty, h_*]} w.$$

For $x \in (h, h_*)$, we have $U_{\bar{X}}^q W_*(x) \leq 0$, but $\mathbf{1}_{(-\infty, h_*]} w(x) > 0$, a contradiction. Thus, (5.12) holds, and the proof is complete. $\square$

5.3. In the case of the put-option on a stock that pays no dividends, we have $q = r$, and the straightforward application of the above scheme faces evident difficulty since $K - e^x$ cannot be represented as the EPV of any stream of payoffs. Indeed, e^x is an eigenfunction of $q - L$: $(q - L)e^x = 0$. Since the put option will not be exercised if $K - e^x$ is negative, we can try to overcome this difficulty by choosing a sufficiently smooth G which coincides with $K - e^x$ on $(-\infty, \log K]$, is negative on $(\log K, +\infty)$, and does not grow (in absolute value) too fast as $x \rightarrow +\infty$, so that $g = (q - L)G$ satisfies $G = U_{\bar{X}}^q g$. However, in this case we need to ensure that $w = U_{\bar{X}}^q g = (qU_{\bar{X}}^q)^{-1}G$ changes sign only once, and $(q - L)qU_{\bar{X}}^q \mathbf{1}_{(-\infty, h_*]} w$ is monotone. Instead, we reformulate the conditions of Theorem 5.2 in terms of the payoff function G and $w = (qU_{\bar{X}}^q)^{-1}G$ so that we avoid the use of a stream g altogether. Notice that $(qU_{\bar{X}}^q)^{-1}G$ is easily calculated if G is a linear combination of exponential functions since $(qU_{\bar{X}}^q)^{-1}e^{zx} = \kappa_q^-(z)^{-1}e^{zx}$. In the proof of Theorem 5.2, the following conditions are essential (and even these conditions can be relaxed):

$$(5.13) \quad (q - L)G \text{ is non-increasing on } (-\infty, h_*),$$

and

$$(5.14) \quad w = (qU_{\bar{X}}^q)^{-1}G \text{ is non-increasing on } (h_*, +\infty).$$

Set $V_*(x) = qU_{\bar{X}}^q \mathbf{1}_{(-\infty, h_*]} w(x)$.

Theorem 5.3. *Let G be continuous, let $w = (qU_{\underline{x}}^q)^{-1}G$ satisfy (5.8), and let (5.13) and (5.14) hold. Then the pair $(\tau_{h_*}^-, V_*(\cdot))$ is the optimal solution to the optimal stopping problem, in the class $\mathcal{M}$.*

Example. For the perpetual American put without dividends, $G(x) = K - e^x$, $w(x) = K - \kappa_q^-(1)^{-1}e^x$, and $(q - L)G(x) = qK$. Clearly, the conditions of Theorem 5.3 are satisfied, and the optimal exercise log-price is $h_* = \ln(K\kappa_q^-(1))$.

5.4. In Boyarchenko and Levendorskii (2000, 2002a,b), we introduced a wide class of Lévy processes: Regular Lévy Processes of Exponential type (RLPE), and, in particular, solved the optimal stopping problem for perpetual American options on a stock driven by an RLPE. A Lévy process is an RLPE if its Lévy density exhibits a regular growth near 0 and exponential decay at infinity. The class of RLPE includes several families of Lévy processes used in empirical studies of financial markets: jump-diffusion processes with exponentially distributed jumps, hyperbolic processes, and extended Koponen's family of Truncated Lévy processes or KoBoL processes (later used under the name CGMY-model in Carr et al. (2002)). For details, see Boyarchenko and Levendorskii (2002a,b,c). Due to certain analytical complications, variance gamma processes were not treated in Boyarchenko and Levendorskii (2000, 2002a,b), although Lévy processes with non-trivial gaussian and variance gamma components were allowed. In Boyarchenko and Levendorskii (2000) we found the exercise boundary for the perpetual American put and rational put price, and in Boyarchenko and Levendorskii (2002a,b), we obtained similar results for calls and put-like and call-like options with more general payoff functions of the form $\max\{G(X_t), 0\}$. For a general G , the optimality of the solution in the class $\mathcal{M}_0$ was shown; the optimality in the class $\mathcal{M}$ was proved for G a linear combination of exponential functions, satisfying certain conditions. The results were formulated in terms of the factors in the Wiener-Hopf factorization formula, or, equivalently, in terms of the supremum and infimum processes, and we conjectured that they held for any Lévy process. Later, for the case of the put and call option, Mordecki (2002) considered arbitrary Lévy process. He showed that the value functions in appropriate discrete time models have the limit as the time step goes to 0, and finished the proof by a sentence which stated that the optimality of the limit "... can be proved exactly as in the discrete time model in Darling et al (1972)"; the precise meaning of the word "exactly" is not clear in this context. It seems unlikely

that the proof in Darling et al (1972), which uses the sufficient conditions in terms of the transition operator of the random walk, can be repeated word by word in the case of a Lévy process, where the verification is typically more involved (see, e.g., Boyarchenko and Levendorskiĭ (2002a, b), and references therein and in Mordecki (2002)). Thus, the proof in Mordecki (2002) is incomplete. Essentially, it is a form of a reasonable guess of the answer, for a particular case of the puts and calls; but Boyarchenko and Levendorskiĭ (2002a, b) obtained the optimal solution in general terms earlier (albeit not for any Lévy process), for wider classes of payoffs.

6. OPTIMAL EXERCISE BOUNDARY AND RATIONAL PRICE OF A PERPETUAL CALL-LIKE OPTION

The statements, arguments, and proofs are mirror reflections of the ones for put-like options. We assume that the supremum process is non-trivial, and g is piece-wise continuous, and satisfies (4.5). Since X satisfies the (ACP)-property, $G = U_X^q g$ is continuous, and it satisfies (4.5). Introduce $w = U_X^q g$, and assume that w is a continuous function that satisfies

$$(6.1) \quad w \text{ changes sign from “} - \text{” to “} + \text{”, and only once.}$$

If g is continuous, w is continuous as well, and a sufficient condition for (6.1) is that g is increasing. This condition makes a perfect economic sense for a perpetual call-like option: the stream of payoffs increases when the stochastic factor increases.

Denote by h^* the solution to the equation

$$(6.2) \quad w(x) = U_X^q g(x) = 0,$$

and set $V^*(x) = qU_X^q \mathbf{1}_{[h^*, +\infty)} w(x)$.

Theorem 6.1. *Let g be non-decreasing, and (6.1) hold. Then the pair $(\tau_{h^*}^+, V^*(\cdot))$ is the optimal solution to the stopping problem, in the class $\mathcal{M}$.*

Example. Assume that $q - \Psi(1) > 0$, and consider the perpetual call option on a dividend paying stock. We have $\kappa_q^+(1) < +\infty$, $G(x) = e^x - K$, $g(x) = (q - L)G(x) = (q - L)e^x - qK$, $w(x) = (qU_X^q)^{-1}G(x) = \kappa_q^+(1)^{-1}e^x - K$. Clearly, g is increasing, condition (6.1) holds, and the optimal exercise log-price is $h^* = \ln(K\kappa_q^+(1))$.

Similarly to Theorem 5.3, we can replace the condition that g is non-decreasing with the following pair of conditions:

$$(6.3) \quad (q - L)G \text{ is non-decreasing on } (h^*, +\infty),$$

and

$$(6.4) \quad (qU_{\underline{X}}^q)^{-1}G \text{ is non-decreasing on } (-\infty, h^*).$$

Theorem 6.2. *Let G be continuous, let $w = (qU_{\underline{X}}^q)^{-1}G$ satisfy (6.1), and let (6.3) and (6.4) hold. Then the pair $(\tau_{h^*}^+, V^*(\cdot))$ is the optimal solution to the stopping problem, in the class $\mathcal{M}$.*

7. INCREMENTAL CAPITAL EXPANSION

Consider a firm whose production function depends only on capital $G = G(K)$. We assume that $G(K)$ differentiable, concave, and satisfies the Inada conditions; the revenue flow is $P_t G(K_t)$, where P_t is the spot price of the firm's output. A similar situation was considered in Dixit and Pindyck (1996) for the geometric Brownian motion model, and extended by Boyarchenko (2004) for geometric Lévy processes. In those papers, the price of the firm's output was modeled as $P_t = e^{X_t}$, where X_t is the Brownian motion and a Lévy process, respectively. In the present paper, we consider more general case, when $P_t = P(X_t)$ is an increasing function of a stochastic factor X_t , which follows a Lévy process. In particular, such a payoff may account for the case when the firm chooses both capital and costlessly adjustable labor as in Abel and Eberly (1999) for the gaussian model, or in Boyarchenko and Levendorskii (2004) for the discrete time model. Should the firm decide to invest a unit of capital, it suffers the installation cost C . At the end of this Section, we allow for a stochastic operational cost as well. In order not to change the notation of the previous Sections, we denote the riskless rate by q . The firm's objective is to choose the optimal investment strategy $\mathcal{K} = \{K_t, t \geq 0\}$, $K_0 = K$, $X_0 = x$, which maximizes the NPV of the firm:

$$(7.1) \quad V(K, x) = \sup_{\mathcal{K}} E^x \left[\int_0^{+\infty} e^{-qt} (P(X_t)G(K_t) - qCK_t) dt \right].$$

To ensure that firm's value (7.1) is bounded, we impose a resource constraint: there exists $\bar{K} < \infty$, such that $K_t \leq \bar{K}$, $\forall t$. Also we assume that X satisfies (2.19), and the function P satisfies (4.5). For the case $P(X_t) = e^{X_t}$, the last two conditions reduce to $q - \Psi(1) > 0$. These conditions and properties of the production function ensure that the value function (7.1) is well defined.

Formally, the manager has to choose both the timing and the size of the capital expansion. However, it is well-known (see, for example, Dixit and Pindyck (1996)) that for each level of the capital stock, it is only necessary to decide when to invest. The manager's problem is equivalent to finding the boundary (the investment threshold),

$h(K; C)$, between two regions in the state variable space (K, x) : inaction and action ones. For all pairs (K, x) belonging to the inaction region, it is optimal to keep the capital stock unchanged. In the action region, investment becomes optimal. To derive the equation for the investment boundary, suppose first that every new investment can be made in chunks of capital, ΔK , only¹. In this case, the firm has to suffer the cost $C\Delta K$, and the EPV of the revenue gain due to this investment can be represented in the form of the EPV of the stream $g(X_t) = (G(K + \Delta K) - G(K))P(X_t) - qC\Delta K$. On the strength of the result of Section 4, the optimal exercise boundary is determined from the equation $U_{\bar{X}}^q g(h) = 0$. In the geometric Lévy case, which we consider first in order to simplify the presentation of the main idea of the proof, the equation for the threshold can be written as

$$(7.2) \quad q^{-1}(G(K + \Delta K) - G(K))\kappa_q^-(1)e^h = C\Delta K.$$

Dividing by ΔK in (7.2) and passing to the limit, we obtain the equation for the optimal threshold, $h^* = h^*(K)$:

$$(7.3) \quad \kappa_q^-(1)G'(K)e^h = qC.$$

Equivalently, the optimal exercise price is

$$(7.4) \quad e^{h^*} = e^{h^*(K)} = \frac{qC}{\kappa_q^-(1)G'(K)}.$$

For the rigorous justification of this limiting argument, see Boyarchenko (2004). Let $h = h(K; \Delta)$ be the solution to (7.2). Then the option value associated with the chunk of capital ΔK , at the price level e^x , is

$$U_{\bar{X}}^q \mathbf{1}_{[h, +\infty)}(x)((G(K + \Delta K) - G(K))\kappa_q^-(1)e^x - qC\Delta K).$$

As $\Delta K \rightarrow 0$, we have $h = h(K; \Delta) \rightarrow h^*(K)$; therefore, dividing by ΔK and passing to the limit, we obtain the formula for the derivative of the option value of future investment opportunities w.r.t. K :

$$(7.5) \quad V_K^{\text{opt}}(K, x) = U_{\bar{X}}^q \mathbf{1}_{[h^*, +\infty)}(x)(G'(K)\kappa_q^-(1)e^x - qC).$$

Substituting C from (7.4) into (7.5) and using the definition of $\kappa_q^-(1)$, we obtain

$$\begin{aligned} V_K^{\text{opt}}(K, x) &= qE \left[\int_0^\infty e^{-qt} G'(K) e^{\bar{X}_t} dt \mid X_0 = 0 \right] \\ &\quad \times E^x \left[\int_0^\infty e^{-qt} (e^{\bar{X}_t} - e^{h^*})_+ dt \right]. \end{aligned}$$

¹The authors are indebted for this simplifying trick to Mike Harrison; the initial proof in Boyarchenko (2004) was more involved.

The last formula factors out the contributions of the infimum and supremum price processes to the marginal option value of capital. The first expectation on the RHS decreases if the probability of downward jumps in prices increases, and the second expectation increases if the probability of positive jumps in prices increases. Hence the marginal option value of capital increases in downward uncertainty and decreases in upward uncertainty. The overall effect of uncertainty is ambiguous.

Notice that the proof of (7.3) in Boyarchenko (2004) was based on the reduction to the case of the perpetual American call, and therefore the generalization for more general dependence on the stochastic factor was not possible. Here the result holds for any continuous increasing revenue flow $R(K, x)$, and the formula for the optimal investment threshold obtains in the form:

$$(7.6) \quad U_{\underline{X}}^q R_K(K, h) = C,$$

where the EPV-operator $U_{\underline{X}}^q$ acts w.r.t. the second argument. For instance, if the firm faces the operational cost $a + bKe^{X_t/2}$, then the revenue flow is $R(K_t, X_t) = e^{X_t}G(K_t) - a - bKe^{X_t/2}$, and instead of (7.4), we now have

$$(7.7) \quad q^{-1}[\kappa_q^-(1)G'(K)e^h - b\kappa_q^-(1/2)e^{h/2}] - C = 0.$$

The function on the LHS in (7.7) changes sign only once, and therefore the solution to equation (7.7) gives the optimal investment threshold.

One can also consider non-exponential dependence of the price on the stochastic factor.

Equation (7.6) says that it is optimal to increase the capital stock the first time the EPV of the marginal revenue, calculated under the assumption that the underlying stochastic process is replaced by the infimum process, reaches or overshoots the marginal cost of investment. This rule reflects and extends the bad news principle spelled out by Bernanke (1983).

REFERENCES

- [1] Abel, A.B., J.C. Eberly: The effects of irreversibility and uncertainty on capital accumulation, *J. Monet. Econ.* 44, 339-377 (1999)
- [2] Bernanke, B.S.: Irreversibility, uncertainty, and cyclical investment, *Quart. J. Econ.* 98, 85-106 (1983)
- [3] Bertoin, J.: Lévy processes (Cambridge tracts in mathematics, vol. 121). Cambridge: Cambridge University Press 1996.
- [4] Boyarchenko, S.I.: Irreversible decisions and record setting news principles, *Amer. Econ. Rev.* (2004), *Forthcoming*

- [5] Boyarchenko, S.I., Levendorskiĭ, S.Z.: Option Pricing for Truncated Lévy Processes. *International Journal of Theoretical and Applied Finance*. 3:3, 549–552 (2000)
- [6] Boyarchenko, S.I., Levendorskiĭ, S.Z.: Perpetual American Options under Lévy Processes. *SIAM Journal of Control and Optimization*. 40:6, 1663–1696 (2002a)
- [7] Boyarchenko, S.I., Levendorskiĭ, S.Z.: Non-Gaussian Merton-Black-Scholes theory. Singapore: World Scientific 2002b
- [8] Boyarchenko, S.I., Levendorskiĭ, S.Z.: Barrier options and touch-and-out options under regular Lévy processes of exponential type. *Annals of Applied Probability*. 12:4, 1261–1298 (2002c)
- [9] Boyarchenko, S.I., Levendorskiĭ, S.Z.: Practical guide to real options in discrete time. Submitted to *J. Econ. Theory* (2004)
- [10] Carr, P., Geman, H., Madan, D.B., and Yor, M.: The fine structure of asset returns: an empirical investigation. *Journal of Business*. 75, 305–332 (2002)
- [11] Darling, D. A., Liggett, T., and Taylor, H. M.: Optimal stopping for partial sums. *Annals of Mathem. Statistics*, 43, 1363–1368 (1972)
- [12] Dixit, A.K., R.S. Pindyck: *Investment under Uncertainty*. Princeton Univ. Press, Princeton, NJ, 1996
- [13] Hilberink, B., and Rogers, L.C.G.: Optimal capital structure and endogenous default. *Finance and Stochastics*, 6, 237–263 (2002)
- [14] Mc Kean, H. P., Jr.: Appendix: A free boundary problem for the heat equation arising from a problem in Mathematical Economics. *Indust. Mang. Rev.*, 6, 32–39 (1965)
- [15] Levendorskiĭ, S.: Pricing of American put under Lévy processes. *International Journal of Theoretical and Applied Finance* (2004)
- [16] Merton, R. C.: The theory of rational option pricing. *Bell Journ. of Economics*, 4, 141–183 (1973)
- [17] Mordecki, E.: Optimal stopping and perpetual options under Lévy processes. *Finance and Stochastics*, 6, 473–493 (2002)
- [18] Sato, K.: *Lévy processes and infinitely divisible distributions*. Cambridge: Cambridge University Press 1999